

**Saccardo, Pier Andrea**

*Sylloge fungorum [...] Vol. III... (1884)*

**Índice**

1. Cd - Cubierta delantera
2. 317
3. 318

# SYLLOGE FUNGORUM

OMNIUM HUCUSQUE COGNITORUM.

DIGESSIT

P. A. SACCARDO

Vol. III.

---

SPHÆROPSIDEÆ et MELANCONIEÆ

---

Fol. 54 — Pretium it lib. (*francs*) 54 .

PATAVII

XV DECEMBRIS MDCCCLXXXIV

SUMPTIBUS AUCTORIS

TYPIS SEMINARII

Iterum impressum

apud

R. Friedländer & Sohn,  
Berlin

R. 39 x 3

ritheciis sparsis, innatis, suberumpentibus, globulosis,  $\frac{1}{5}$  mill. d., ostiolo brevi obtuso; sporulis globuloso-ellipsoideis, 3,6-4,5  $\mu$ . d., fuligineis, uniguttatis.

*Hab.* in caulibus *Euphorbiæ silvaticæ*, Elbeuf in Gallia (MALBRANCHE).

64. **Coniothyrium Parietariæ** Sacc. et Speg. Mich. I. p. 204. — Peritheciis cortice secedente superficialibus, subhemisphæricis, 200-300  $\mu$ . diam., umbilicatis, carbonaceis; sporulis ovoideo-globulosis, 4-5  $\approx$  2-3  $\frac{1}{2}$ , lutescenti-olivaceis.

*Hab.* in radicibus emortuis *Parietariæ officinalis*, Conegliano Italiæ bor. (SPEGAZZINI).

65. **Coniothyrium herbarum** Schulz. et Sacc. Micr. Slav. n. 26, Schulz. 1787 Ill. Fung. Slav. n. 271. — Peritheciis subgregariis, initio epidermide velatis, lenticularibus,  $\frac{1}{10}$ - $\frac{1}{6}$  mill. diam., poro pertusis; sporulis oblongo-teretiusculis, utrinque rotundatis, 5  $\approx$  2, brunneolis.

*Hab.* in caulibus *Asclepiadis Syriacæ* prope Vinkovce in Slavonia. — Præcipue peritheciis lenticulari-applanatis dignoscenda species.

66. **Coniothyrium æquivocum** (Ces.) Sacc. *Phoma æquivocum* Ces. 1788 Myc. Born. p. 25. — Peritheciis subcutaneis, in caule exsiccato, sparsis, globoso- vel ellipsoideo-depressis, ostiolo papillari emergente; sporulis ovalibus obtusis, fuscis, 3  $\approx$  2.

*Hab.* in caulibus exsiccatis, Sarawak Bornei (BECCARI).

### III. In Monocotyledoneis.

67. **Coniothyrium concentricum** (Desm.) Sacc. Mich. I. p. 204, *Phoma concentrica* Desm. Ann. Sc. Nat. 1840, XIII. p. 189. — Peritheciis lentiformibus, subastomis, contextu parenchymatico fuligineo; sporulis e globoso ovoideis, 4-5  $\approx$  3-4, initio hyalinis, dein flavidis, tandem fuligineis, plerumque 1-guttulatis.

*Hab.* in foliis *Yuccæ gloriosæ*, Yuc. filamentosæ, Yuc. aloifoliæ, Fourcroyæ excelsæ in Italia, Gallia, Belgio, Lusitania, Britannia et Amer. bor. — Perithecia non semper concentrica disposita, modo epidermide nigricante nunc immutata velata.

— Var. **Agaves**: peritheciis sparse gregariis, erumpenti-superficialibus, globulosis. In foliis Agaves americanæ in Italia, Lusitania, Austria. — \***C. inops** Sacc. Mich. II. p. 142, *Phoma concentrica* Rav. et Cooke F. A. n. 243. In foliis *Yuccæ filamentosæ*. — A *C. concentrico* typico differt peritheciis minus evolutis,  $\frac{1}{4}$  mill. diam., globoso-conicis, immersis et sporulis, 3-

Matutino

Probabilitate  
C. Agave (Monst) late  
sert.  
Hispania (Hypox)

$3\frac{1}{2}$   $\mu$ . d., globosis, minoribus, olivaceo-fuligineis, 1-guttulatis. Ob perithecia imperfecta forte huc pertinet *Papularia concentrica* Kickx Flandr. II. p. 90, nec *C. concentricum* Auct. — *C. herbarum* Cooke et Ellis Grevill. VI. p. 1 affine est, sed distingui potest.

68. **Coniothyrium Henrquesii** Thüm. Contr. Myc. Lus. n. 338. — 1790

Peritheciis amphigenis, magnis, hemisphærico-emersis vel vix conoideis, primo epidermide tectis, postremo cuticulâ dilaceratâ cinctis, atris, in disco articulato, magno, elevato, gregarie dispositis vel plerumque solitariis et sine ordine; sporulis bacillaribus vel longe ellipticis, rectis vel interdum subcurvulatis, utrinque subobtusatis, simplicibus, fuscis, 7-8  $\mu$ .

*Hab.* in foliis emortuis Fourcroyæ Sellowii in horto botanico Coimbra in Lusitania (MOLLER).

69. **Coniothyrium biforme** Winter Contr. Myc. Lus. n. 790. — Peritheciis 1791

nunc immersis, demum erumpentibus, usque 300  $\mu$ ., nunc superficialibus, 130-140  $\mu$ ., subglobosis, nitidis, atris, membranaceis, sub microscopio atro-violaceis; sporulis angulato-globosis, fuscis, 5-7  $\mu$ . diam.

*Hab.* ad folia emortua Fourcroyæ in horto botanico Coimbra in Lusitania.

70. **Coniothyrium Agaves** (Mont.) Sacc. *Phoma Agaves* Dur. et Mont. 1792

Fl. Alg. p. 605, Mont. Syll. n. 980. — Gregarium, erumpens, punctiforme, inquinans; peritheciis rotundatis, minutissimis, atris, opacis aut nitidis; sporophoris brevibus; sporulis ovato-globosis, continuis, obscure fuliginosis.

*Hab.* in foliis emarcidis Agaves americanæ Alger (DURIEU).

(Agave) — An eadem ac var. *Agaves Coniothyrii concentrici*?

71. **Coniothyrium herbarum** Cooke et Ellis Grev. t. 96, f. 32. — Pe- 1793

ritheciis convexis, brunneis, primo tectis, membranaceis; sporulis subglobosis, brunneis, 4-5  $\mu$ . diam.

*Hab.* in pedunculis Yuccæ filamentosæ.

72. **Coniothyrium Palmarum** Corda Icon. Fung. IV. p. 38, t. 8, f. 106. 1794

— Gregarium; peritheciis oblongis, convexis, plicatulis, atris, nitentibus, maculâ fusca, margine saepe rubescente insidentibus; sporulis ellipsoideis in sicco coffeæformibus, semipellucidis, 8-10  $\times$  4. crasse 2-nucleatis, dein obscure fuligineis.

*Hab.* in foliis languidis vel emortuis Chamæropis humilis in Italia australi, in Sicilia nec non in foliis Phœnicis dactyliferae in horto botanico Coimbra in Lusitania.

73. **Coniothyrium borbonicum** Thüm. Contr. Myc. Lusit. n. 339. — 1795